



INTERNATIONAL JOURNAL OF PHARMACEUTICAL RESEARCH AND DEVELOPMENT (IJPRD)

Platform for Pharmaceutical Researches & Innovative Ideas
www.ijprd.com

ANTIMICROBIAL ACTIVITY AND CYTOTOXICITY OF SEED EXTRACTS OF *MESUA NAGASSARIUM* (BURM.F.)

Md. Masud Pervez^{*1},
Md. Arifur Rahman¹, Md. Khosruzzaman Molla¹, and Md Al Amin Sikder¹

¹Department of Pharmacy, State University of Bangladesh, Dhaka-1205, Bangladesh

ABSTRACT

In the present study the in vitro antimicrobial activity, along with minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC), of different extracts of Seeds of *Mesua nagassarium* were evaluated against 13 pathogenic microorganisms. The methanol extract and its pet-ether and carbon tetrachloride soluble fractions showed the highest antimicrobial activity. The carbon tetrachloride soluble fraction showed the maximum inhibition zone of 22.33 mm against *Bacillus megaterium* with MIC and MBC values of 7.90 µg/ml and 250 µg/ml, respectively. Ciprofloxacin (30 µg/disc) was used as standard antimicrobial agent. In the *Artemia salina* lethality bioassay, the crude methanol extract and its carbon tetrachloride soluble fraction showed significant cytotoxicity with LC50 of 3.14 and 1.74 µg/ml, respectively as compared vincristine sulphate (LC50 value 0.543 µg/ml).

Correspondence to Author



Md. Masud Pervez

Department of Pharmacy, State
University of Bangladesh, Dhaka-
1205, Bangladesh

Email

Mparvezph10@gmail.com

Key Words

Mesua nagassarium,
Antimicrobial Activity,
Cytotoxicity, skin eruptions,
dandruff, etc.

INTRODUCTION

Mesua nagassarium (Bengali - nagesar, nageswar) is a medium-sized or fairly large evergreen tree up to 36 m tall. A mixture of pounded kernels and seed oil isolated from this plant is used for poultice as wounds. The seed-oil is used for treating itch and other skin eruptions, dandruff and rheumatism (Orwa et al., 2009). A mixture of pounded kernels and seed oil isolated from this plant is used for poultice as wounds. The seed-oil is used for treating itch and other skin eruptions, dandruff and rheumatism. The flowers are known to be useful for the treatment of severe colds, bleeding haemorrhoids, dysentery with mucus, excessive thirst, excessive perspiration, cough and digestion, rheumatism and iron induced lipid peroxidation (Yadav et al., 2010; Konwarh et al., 2010). Pet ether extracts of leaf of *M. nagassarium* also shows highly antibacterial activity (Shikder et al., 2011). Phenolic extract of seed oil of *M. nagassarium* revealed potent antiasthmatic effect (Bhide, 1977). As a part of our ongoing program to investigate the unexplored bioactivity of traditionally used medicinal plant of Bangladesh we studied the antimicrobial and cytotoxicity of seeds of *M. nagassarium* were evaluated.

MATERIALS AND METHODS

Plant Material

The seeds of *Mesua nagassarium* were collected from Dhaka in April 2011. A voucher specimen for this plant has been maintained in Bangladesh National Herbarium, Dhaka, Bangladesh (Accession no. DACB- 35158). The powdered plant sample (600 gm) was soaked in 2.5 L of methanol for 7 days and then filtered through a cotton plug followed by Whatman filter paper number 1. The extract was concentrated with a rotary evaporator at low temperature (40-45 °C) and reduced pressure. The concentrated methanolic extract (MES) was partitioned by modified Kupchan method (Van Wageningen et al., 1993) and subsequent evaporation of solvents yielded pet-ether (PESF), carbon tetrachloride (CTCSF), dichloromethane (DCMSF) and aqueous (AQSF) soluble fractions which were used for the experimental processes.

Antimicrobial screening

The disc diffusion method (Bauer et al., 1966, Rahman and Rashid, 2008) was used to test antimicrobial activity of the extractives against thirteen bacteria (Table-1), collected as pure cultures from the Institute of Nutrition and Food Sciences (INFS), University of Dhaka, Bangladesh. Solutions of known concentration (400 µg/ml) of the test samples were made by dissolving measured amount of the samples in calculated volume of solvents. Dried and sterilized filter paper discs (6 mm diameter) were then impregnated with known amounts of the test substances using micropipette and the residual solvents were completely evaporated. Discs containing the test materials (400 µg/disc according to disc diffusion method (Bauer et al., 1966, Rahman and Rashid, 2008) were placed on to nutrient agar medium uniformly seeded with the test microorganisms. Standard disc of Ciprofloxacin (30 µg/disc) and blank discs (impregnated with solvents followed by evaporation) were used as positive and negative control, respectively. These plates were then kept at low temperature (4 °C) for 24 hours to allow maximum diffusion of the test materials and Ciprofloxacin. The plates were finally incubated at 37 °C for 24 hours to allow maximum growth of the organisms. The test material having antimicrobial activity inhibited the growth of the microorganisms and a clear, distinct zone of inhibition was visualized surrounding the discs. The antimicrobial activity of the test agents was determined by measuring the diameter of zone of inhibition expressed in mm. The experiment was carried out in triplicate and the mean value was taken.

Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) determination

Minimum inhibitory concentrations are important to monitor the activity of new antimicrobial agents (Jennifer, 2001) and is generally regarded as the most basic laboratory measurement of the activity of an antimicrobial agent against an organism (Turnidge et al., 2003). The MIC was determined for MES, PESF and CTCSF extractives by serial tube dilution technique (Jennifer, 2001). Thirteen test tubes were taken, ten of which were marked as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 and the rest three were assigned as T_M (only medium), T_{MC} (Medium +extractive solution) and T_{MI} (medium +

inoculum). In all the test tubes containing the calculated amount of broth media, the sample from the mother solution was added with serial dilution which gave different sample solution having concentration ranged from 1000 µg/ml to 0 µg/ml. Lastly the inoculum was added to the test tubes, shaken using rotamixer and incubated at 37 °C for 24 hours. The control test tube T_M, containing the medium only was used to ascertain the sterility of the medium. After 24 hours the test tubes were checked for the microbial growth and the clear test tubes compared to McFarland turbidity standards were marked for the MIC determination. Serial dilution of the liquid medium and the sample is inoculated with inoculum whose turbidity was compared with 0.5 Mcfarland standards. The lowest concentration (highest dilution) of antibiotic preventing appearance of turbidity is considered to be the minimal inhibitory concentration (MIC). At this dilution the antibiotic is bacteriostatic. The minimum bactericidal concentration (MBC) was determined by sub-culturing the broth media into fresh agar media from each tube in which no growth was visible. The growth of one colony indicated a 99.8% fall in viable count (Brumitt et al., 1984).

Brine shrimp lethality bioassay

Brine shrimp lethality bioassay (Meyer et al., 1982, McLaughlin et al., 1998) technique was applied for the determination of general toxic property of the plant extractives. DMSO solutions of the samples were applied against *Artemia salina* in a 1-day *in vivo* assay. For the experiment, 4 mg of each of the methanol extract as well as pet-ether, carbon tetrachloride, chloroform and aqueous soluble fractions of methanol extract were dissolved in DMSO and solutions of varying concentrations (400, 200, 100, 50, 25, 12.50, 6.25, 3.125, 1.563, 0.781 µg/ml) were obtained by serial dilution technique using DMSO. Vincristine sulphate (10 µg/ml, 5 µg/ml, 2.5µg/ml, 1.25 µg/ml, 0.625 µg/ml, 0.3125 µg/ml, 0.15625 µg/ml, 0.078125 µg/ml, 0.0390 µg/ml) was used as positive control.

Results and Discussion

The crude methanolic extract of leaves of *M. nagassarium* (MES) and its pet-ether (PESF), carbon tetrachloride (CTCSF), dichloromethane (DCMSF) and aqueous (AQSF) soluble partitionates were subjected to antimicrobial screening at 400 µg/disc. Among the extractives, MES, PESF and CTCSF exhibited very strong antimicrobial activity (Table 1).

Table 1: *In vitro* antibacterial activity of the extracts from *M. nagassarium* seeds

Test microorganisms	Diameter of zone of inhibition (mm)					
	MES	PESF	CTCSF	DCMSF	AQSE	Ciprofloxacin
Gram positive bacteria						
<i>Bacillus cereus</i>	21.3±0.57	21.3±2.08	22.33±2.08	10.7±1.15	10.3±2.08	40.33±3.78
<i>B. megaterium</i>	23.0±1.0	20.0±1.73	22.33±1.53	12.0±1.73	8.67±0.58	44.66±1.52
<i>B. subtilis</i>	20.3±1.52	21.0±1.73	22.67±1.15	12.3±2.08	10.3±0.58	40.66±2.08
<i>Staphylococcus aureus</i>	20.3±1.52	20.7±1.15	22.0±1.0	10.3±0.57	9.67±0.58	43.0±1.73
<i>Sarcina lutea</i>	21.0±1.0	21.3±3.51	21.67±2.08	11.0±1.0	11.3±1.53	41.0±2.0
Gram negative bacteria						
<i>Escherichia coli</i>	20.0±1.73	20.3±0.58	21.33±1.15	10.0±1.0	9.67±0.57	44.67±1.53
<i>Pseudomonas aeruginosa</i>	21.3±1.52	20.0±0	20.0±2.0	10.3±0.57	11.0±1.73	44.0±2.65
<i>Salmonella paratyphi</i>	21.7±2.08	22.3±2.08	22.67±2.52	9.67±0.57	10.7±2.08	44.33±2.08
<i>S. typhi</i>	17.20±2.08	19.3±1.15	19.03±3.06	9.67±0.57	12.3±2.08	43.66±1.52

<i>Shigella boydii</i>	22.7±2.51	20.7±3.07	22.0±2.65	9.67±0.57	11.7±1.53	44.33±2.08
<i>S. dysenteriae</i>	20.7±1.15	20.7±1.53	19.67±2.08	10.0±2.0	10.7±3.06	41.33±1.52
<i>Vibrio mimicus</i>	21.3±1.15	22.3±2.31	23.0±1.0	11.0±1.73	11.7±2.08	43.33±2.88
<i>V. parahemolyticus</i>	21.7±2.08	22.7±2.52	22.67±2.31	11.0±1.0	12.0±2.0	44.0±1.73

Values are expressed as mean ± S.D. (n=3)

MES= Methanol extract, PESF= Pet-ether soluble fraction, CTCSF= Carbon tetrachloride soluble fraction, DCMSF= dichloromethane soluble fraction, AQSF= aqueous soluble fraction.

The carbon tetrachloride soluble fraction (CTCSF) revealed the highest inhibition against microbial growth having zone of inhibition ranged from 17.20 mm to 22.33 mm. The maximum zone of inhibition produced by CTCSF was found to be 22.33 mm against *B. megaterium*. This partitionate also showed significant antimicrobial growth having zone of inhibition 22.67 mm against *S. paratyphi* (Table 1). The methanol extract also demonstrated significant inhibition of microbial growth having zone of inhibition ranging from 17.20 mm to 23.0 mm. This extract exerted highest inhibitory activity against *B. megaterium* (having zone of inhibition of 23.0 mm). Relatively less polar compounds revealed better antimicrobial activity as the pet-ether and carbon tetrachloride soluble partitionates showed significant antimicrobial activity than the aqueous and chloroform soluble partitionates of methanol extract.

As MES, PESF, CTCSF showed better inhibitory activity against the microorganisms, the minimum inhibitory concentration (MIC) required to inhibit the growth of organisms were measured in this study. The MIC value of methanol extract (where no bacterial growth was observed in broth media) was found to be 7.90 µg/ml and the MBC value (where no bacterial growth was observed in agar media) was found to be 250 µg/ml against *B. megaterium* (Table 2). No microbial growth was observed in the test tubes T_M (containing medium only) and T_{MC} (medium + test sample, no inoculum) indicating that the medium and the test sample were not contaminated by microorganism and the total investigation was performed properly in the sterile condition. Microbial growth was observed in the test tube T_{MI} (medium + inoculums) revealing the fact that there was no problem with the subcultured microorganisms.

Table 2: Determination of MIC and MBC (µg/ml) of the extracts from *M. nagassarium* seeds

		Determination of MIC and MBC (µg/ml)							
		MES		PESF		CTCSF		Ciprofloxacin	
Test microorganisms		MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
Gram positive bacteria									
<i>Bacillus megaterium</i>		7.90	250	15.62	250	7.81	250	1.25	2.5
<i>Bacillus subtilis</i>		31.25	500	15.62	250	62.5	500	0.62	1.25
<i>Staphylococcus aureus</i>		31.25	500	15.62	500	7.81	250	0.31	0.62
<i>Sarcina lutea</i>		15.62	500	15.62	500	7.81	250	1.25	5.0
Gram negative bacteria									
<i>Escherichia coli</i>		62.5	1000	31.25	500	31.25	250	1.25	2.5
<i>Pseudomonas aeruginosa</i>		31.25	500	62.5	500	15.62	500	0.62	2.5

<i>Salmonella typhi</i>	62.5	1000	31.25	500	31.25	500	0.31	1.25
<i>Shigella dysenteriae</i>	31.25	500	31.25	250	62.5	500	0.31	2.5
<i>Vibrio parahemolyticus</i>	31.25	1000	15.62	500	15.62	500	0.62	2.5

MIC = Minimum inhibitory concentration, MBC= Minimum bactericidal concentration.

In case of brine shrimp lethality bioassay, the lethality of the methanol extract (MES) and its pet-ether (PESF), carbon tetrachloride (CTCSF), dichloromethane (DCMSF) and aqueous (AQSF) soluble fractions were evaluated against *A. salina*. (Table 3) shows the results of the brine shrimp lethality testing after 24 hours of exposure to the

samples and the positive control, vincristine sulphate (VS). The carbon tetrachloride soluble fraction (CTCSF) showed potent cytotoxic activity having LC₅₀ of 1.94 µg/ml as compared to 0.453 µg/ml for vincristine sulphate.

Table 3: Cytotoxic activity of different partitionates of *M. nagassarium*.

Sample	Cytotoxic activity (LC ₅₀ mg/ml)
VS	0.453±0.002
MES	3.14±0.188
PESF	4.83±0.025
CTCSF	1.94±0.11
DCMSF	12.64±1.03
AQSF	7.86±0.09

The average values of three calculations are presented as mean ± S.D. (standard deviation); VS = Vincristine sulphate; MES = Methanolic extract; PESF = Pet-ether soluble fraction; CTCSF = Carbon tetrachloride soluble fraction; DCMSF= dichloromethane soluble fraction; AQSF = Aqueous soluble fraction of the methanolic extract of *M. nagassarium*

CONCLUSIONS

The results of *in vitro* microbial screening of *M. nagassarium* indicated that the methanol extract and its pet-ether (PESF) and carbon tetrachloride (CTCSF) soluble partitionates have strong antimicrobial activity.

REFERENCES

1. Bauer AW, Kirby WMM, Sherris JC, Tuck M. 1966. Antibiotic susceptibility testing by a standardized disc diffusion method. J Am clin Pathol 45: 493 - 496.
2. Bhide MB. 1977. Studies on the antiasthmatic activity of *Mesua ferrea*. Haff. Inst. 5: 27.
3. Brumitt W, Hamilton-Miller MT, Gooding A. 1984. Importance of methodology in
4. determining bactericidal and bacteriostatic activities of Azlocillin and ticarcillin against *Pseudomonas aeruginosa*. J. Med Microbiol 17: 37 - 44.
5. Jennifer MA. 2001. Determination of minimum inhibitory concentrations. J. Antimic. Chemother 48: 5 - 16.
6. Konwarh R, Kalita D, Mahanta C, Mandal M, Karak N. 2010. Magnetically recyclable, antimicrobial, and catalytically enhanced polymer-assisted "green" nanosystem-immobilized *Aspergillus niger* amyloglucosidase. Appl Microbiol Biotechnol 87: 1983 - 1992.
7. McLaughlin JL, Rogers LL. 1998. The use of biological assays to evaluate botanicals. Drug Information J 32: 513 - 524.

8. Md. Al Amin SIKDER, Mohammad A. KAISAR, Md. Masud PARVEZ, A.K.M. Nawshad HOSSIAN, Farzana AKHTER & Mohammad A. RASHID 2011. Preliminary Antimicrobial Activity and Cytotoxicity of Leaf Extracts of *Mesua nagassarium* (Burm.f.). 83-87.
9. Meyer BN, Ferrigni NR, Putnam JE, Jacobsen JB, Nicholsand DE, Mclaughlin JL. 1982. Brine shrimp; a convenient general bioassay for active plant constituents. *Planta Med* 45: 31 - 34.
10. Orwa C, Mutua A, Kindt R, Jamnadass R, Simons A. 2009. *Agroforestry Database*. 4.0.
11. Rahman MS, Rashid MA. 2008. Antimicrobial activity and cytotoxicity of *Eclipta prostrata*. *Oriental Pharm Exp Med* 8: 47 - 52.
12. Turnidge JD, Ferraro MJ, Jorgensen JH. 2003. *Manual of Clinical Microbiology*. 8th Ed. Washington. American Society of Clinical Microbiology. p. 1103.
13. Van Wagenen BC, Larsen R, Cardellina JH, Ranzazzo D, Lidert ZC & Swithenbank C. 1993. Ulosantoin, a potent insecticide from the sponge *Ulosa ruetzleri*. *J Org Chem* 58: 335 - 337.
14. Yadav AS, Bhatnagar D. 2010. Inhibition of iron induced lipid peroxidation and antioxidant activity of Indian spices and Acacia in vitro. *Plant foods Hum Nutr Mar* 65: 18 – 24.
