



**COMPARATIVE ANALYSIS OF METALLIC ELEMENTS AMONG MENOPAUSAL, PREGNANT AND UN MARRIED
FEMALES BY USING FINGERNAILS****Aima Iram Batool^{1*},**Naima Huma Naveed¹, Fayyaz ur Rehman², Aamir Ali¹, Nabiha Riaz¹, Saima Farah Naz¹ and Tasawar Ali¹¹Department of Biological Sciences University of Sargodha, Punjab, Pakistan²Department of Chemistry University of Sargodha, Punjab, Pakistan**ABSTRACT**

Nails being noninvasive material are good biomarkers of metallic elemental status in human body. Three categories of females viz unmarried, pregnant and menopause were selected for present study. Excess or deficiency of metallic elements among unmarried, pregnant and menopause produce differential effects. Among unmarried ones leads towards menstrual disorders. Pregnant ones are more susceptible in terms of fetus safety, among which they can produce adverse reproductive outcomes. Menopausal can suffer from health disorders due to imbalance of these metals. Knowledge about metallic elements in females body is important in this regard because they are part of whole body machinery and there deficiency, excess or any imbalance leads towards different disorders The metals under consideration were lead, iron, cadmium, chromium, manganese, copper and zinc. Concentration of lead, cadmium and manganese was significantly elevated among menopausal females matched to unmarried and pregnant females. Iron level and copper level were higher among un-married as compared to other two categories. Pregnant females had elevated level of zinc and chromium.

Correspondence to Author**Aima Iram Batool**Department of Biological Sciences
University of Sargodha,
Punjab, Pakistan**Email:** aima_uos@yahoo.com**INTRODUCTION**

Metal determination in human tissues is the most common application of biological monitoring for screening and diagnosis of metal exposures. Nails as biopsy materials are suggested as more attractive biomarkers in assessing heavy metals environmental exposure. They are preferred biological medium because of ease of collection, Available online on www.ijprd.com

storage, convenience, their usefulness in estimating intake of minerals in nutritional studies and ease of handling (Daniel *et al.*, 2004)

Metal contamination issues are becoming increasingly common because of Industrialization, urbanization, mining operations, increased vehicular traffic, use of fertilizers and pesticides in agriculture. The essential trace elements getting

displaced from the metabolic active sites, by accumulation of toxic trace elements from the working environment. For example chromium is an essential trace element required for normal carbohydrate, lipid, and protein metabolism (Vincent,2004) and normalizes blood sugar levels (Anderson,1980). Chromium can be toxic at higher levels, as found in industrial exposure, causing hepatic per oxidative damage while chromium deficiency depresses nucleic acid synthesis and decreases fertility, also results in impaired glucose tolerance, hyperglycemia, and glycosuria that cannot be controlled with insulin (Anderson,1998; Jeejeebhoy,1999).

The involvement of copper in so many essential biochemical reactions in humans has long been recognized. Copper is required as a cofactor for many enzymes such as cytochrome oxidase (Friberg *et al.*,1986). Copper poisoning frequently associated with suicidal intent include: nausea, vomiting, diarrhea, hypotension, jaundice, hematuria, anuria, coma and death (Chuttani *et al.*,1965). Copper deficiency leads to anaemia defective wool keratinization, abnormal bone formation, arterial and cardiac aneurysm (Hart *et al.*,1928; Shields *et al.*,1965).

Iron cation is crucial for the delivery of oxygen within the body through interactions with hemoglobin and myoglobin. Iron is also the main part of cytochrome C which is responsible for electron transfer within mitochondria. Therefore, a high iron concentration may have an indirect impact on fatty acid oxidation and ATP production in mitochondria (Griffith,1995; Clement,1994). Iron deficiency is common in overweight and obese subjects (Nead *et al.*,2004). Fe deficiency is involved in the pathophysiology of restless legs syndrome (Tan *et al.*,2006) also associated with alterations in many metabolic processes that affects brain functioning, among whom are neurotransmitter metabolism, protein synthesis, organogenesis (Beard,1999).

Present investigation study was carried out in order to determine the concentration of different metals in the nails samples of females, resident of Sargodha district which was categorized on the Available online on www.ijprd.com

basis of age and physical condition i.e. pregnant, un married and menopausal.

MATERIAL AND METHOD

Introduction of the study site

Sargodha is the city of Sargodha District, Punjab, Pakistan. It is located in the north-east of Pakistan. It is the eleventh largest city of Pakistan and is known as Pakistan's best citrus-producing area. It is an agricultural trade center with various industries. Pakistan's largest Airbase is situated in Sargodha and known as 'city of eagles'.

Collection and Analysis of Data:

The questionnaires were filled by face to face conversation regarding marital status, no of children. After it 75 females were selected.

Investigation study was conducted on female individuals residing in Sargodha district. The population under study was categorized on the basis of age and physical condition like unmarried, pregnant and menopause. 25 females who all were healthy with age ranging from 20-35 were included in unmarried group while 25 females who were 3-7 months pregnant were categorized in pregnant group. Menopausal group was consisted of 25 menopausal female (age 40 – 70) all were healthy volunteers.

A weighed quantity of (100 mg) of nails sample was taken in properly washed and labelled Erlenmeyer flask. Acid digestion was carried out by using nitric acid and hydrogen per oxide (Rodushkin and Axelsson,2000). 4ml of nitric acid was added to each sample and incubated it overnight at room temperature. The samples were heated on hot plate, cooled and then 2ml of perchloric acid was added to them. Then volume up to 25ml was made up with the help of doubled distilled water. After filtering solution was transferred into marked Teflon bottles. After wet acid digestion samples of nails were run through Atomic absorption spectrophotometer (AA 6600 Shimadzu) for determination of Pb, Zn, Fe, Cu, Cr, Cd and Mn.

RESULT AND DISCUSSION

The present investigation exhibits a great variation in metal concentration among pregnant, un married and menopausal females due to some natural factors i.e. climate change and demographic factors. The metals under consideration were lead, iron, cadmium, chromium, manganese, copper and zinc

Cadmium is considered as a toxic heavy metal with no known metabolic function in the body (Lauwerys *et al.*,1994). Cd concentration in nails of females was 0.0042-0.0130 $\mu\text{g/g}$ and highest values were present among Menopause ones (0.0067-0.00112 $\mu\text{g/g}$) (fig1). Were *et al.*,2008 reported Cd concentration in nails as $0.44\pm 0.06 \mu\text{g/g}$ and $0.73\pm 0.08 \mu\text{g/g}$ while according to Mehra and Juneja(2003) the Cd level was $0.321\pm 0.21 \mu\text{g/g}$ that all are higher than ours. Cd nail concentration in Pregnant ranged from 0.0042-0.0130 $\mu\text{g/g}$ while in Un-married it was observed as 0.00128-0.00212 $\mu\text{g/g}$.

Lead is naturally occurring metal that people have used almost since the beginning of

civilization. Concentration of lead in nails samples was recorded as 14.16 $\mu\text{g/g}$ - 0.2 $\mu\text{g/g}$ that are in agreement with Mehra and Juneja,2003who reported its level as $7.60\pm 6.44 \mu\text{g/g}$ and the highest value of Pb nail concentration was found in menopause females (8.248-14.16 $\mu\text{g/g}$) while Pb was detected as 1.213-3.48 $\mu\text{g/g}$ for unmarried and 0.21-3.21 $\mu\text{g/g}$ for pregnant ones.


Iron cation is crucial for the delivery of oxygen within the body through interactions with hemoglobin and myoglobin. High iron concentration may have an indirect impact on fatty acid oxidation and ATP production in mitochondria (Griffith,1995; Clement,1994).Mehra and Juneja (2003) reported Fe concentration in nails as $147.18\pm 57.65 \mu\text{g/g}$ that does not substantiate with our ones (0.304-2.611 $\mu\text{g/g}$). Fe concentration was found highest among Un-married in range of 2.27-3.116 $\mu\text{g/g}$ while Fe concentration ranged from 0.304 $\mu\text{g/g}$ to 1.545 $\mu\text{g/g}$ in Pregnant and from 1.531 to 2.61 $\mu\text{g/g}$ in Menopausal females.

Minerals	Source of variation	Df	MS	F value
Lead	Un married× pregnant× menopause	2	763.0545	544.5965*
Iron	Un married × pregnant× menopause	2	17.35626	195.456*
Cadmium	Un married× pregnant × menopause	2	0.000317	2.17E-28*
Chromium	Un married × pregnant × menopause	2	0.522911	313.6482*
Manganese	Un married× pregnant× menopause	2	0.206346	17.97309*
Copper	Un married× pregnant× menopause	2	144.2705	2.000352*
Zinc	Un married× pregnant× menopause	2	143.4668	1.016217*

Chromium, in the form of naturally occurring dinicotinic acid-glutathione complex, also known as glucose tolerance factor (GTF), is vital for carbohydrate metabolism as it potentiates the action of insulin (Merz,1982). Statistically significant difference was found among un married

vs pregnant vs menopausal females. Cr nail concentration ranged from 0.522-0.3040 $\mu\text{g/g}$, 0.4844-0.5109 $\mu\text{g/g}$ and 0.3675-0.4276 $\mu\text{g/g}$ of Un-married, Pregnant and Menopause respectively. Were *et al.*, 2008worked on nails toxicity and their

finding are as 0.73 $\mu\text{g/g}$, 0.44 $\mu\text{g/g}$ that is in agreement with our observation.



Manganese is recognized as playing an important role in the functioning of isocitric dehydrogenase, an important control enzyme in the regulation of Krebs Cycle (Harper,1979). In case of nails Mn concentration evaluated by Mehra and Juneja,2003 values were as 3.16, 3.81 $\mu\text{g/g}$ and 1.31 $\mu\text{g/g}$, while our are in the range of 0.054 – 0.2231 $\mu\text{g/g}$ among un married that are lower than that of Mehra and Juneja. The highest peak observed in Menopause for Mn nail concentration was (0.2406-0.3102 $\mu\text{g/g}$) while 0.1489-0.2231 $\mu\text{g/g}$ concentration was recorded for pregnant ones.

The involvement of copper in so many essential biochemical reactions in humans has long been recognized. Copper is required as a cofactor for many enzymes such as cytochrome oxidase (Friberg *et al.*,1986). Cu nails concentration ranges from 2.955-31.645 $\mu\text{g/g}$, 7.575-10.705 $\mu\text{g/g}$ and 7.715-14.04 $\mu\text{g/g}$ in Un-married, Pregnant and Menopause so Pregnant shows the highest peak on graph than others. Mehra and Juneja Mehra and Juneja,2003 reported 8.48 $\mu\text{g/g}$ Cu in nail samples

that is in agreement with our values (7.575 – 14.04 $\mu\text{g/g}$).

Zinc, one of the important essential trace mineral is required in gene expression and the regulation of cellular growth and differentiation. Zn dependent enzymes are involved in macronutrient metabolism and cell replication(Hays and Swenson,1985; Arinola *et al.*,20080. The ranges of Zn nail concentration in Pregnant, Menopause and Un-married were 9.095-68.54 $\mu\text{g/g}$, 6.725-34.89 $\mu\text{g/g}$ and 6.68-51.855 $\mu\text{g/g}$ respectively. Pregnant Zn nail concentration value is maximum in our observations.

REFERENCES

1. Anderson, R.A.1980.Essentiality of chromium in humans. J. Sci. Tot. Environ. 75-86
2. Anderson, R.A. (1998). Chromium, glucose intolerance and diabetes. J. Am. Coll. Nutr., 17:548–555.

3. Arinola, O.G., Nwozo, S.O., Ajiboye, J.A., Oniye, A.H. 2008. Evaluation of trace elements and total antioxidant status in Nigerian cassava processors. *Pak. J. Nutr.* 7(6): 770-772.
4. Beard, J.L. 1999. Iron deficiency and neural development: an update-*Arch. J. Latin. Nutr.* 49: 34-39.
5. Chuttani HK, Gupta PS and Gukati S (1965). Acute copper Sulphate Poisoning. *Am. J. Med.* 39:849.
6. Clement, F. 1994. Regulation of iron balance in human. *Blood*; 84:1697-1702.
7. Daniel, C. R., Piraccini, B. M and Tosti, A. 2004. The nail and hair in forensic science. *J. Am. Acad. Dermatol.* 50(2):258–261.
8. Friberg, L., Norberg, G. F., and Voukvb. 1986. Handbook on toxicology of metal, vol.ii, specific metals. Elsevier science publisher, Amsterdam.
9. Griffith, H. W. 1995. Complete guide to vitamins, minerals and supplements. Fisher book company press, Newyork.
10. Harper, H. A. 1979. Review of physiological chemistry, 17th Ed. Los Altos, Lange medical publications, California.
11. Hart EB, Steenboch H, Waddell J, Elvehjem CA (1928). Iron in nutrition, VII. Copper as a supplement to iron for haemoglobin building in rat. *J. Biol. Chem.*, 77:797.
12. Hays, V.W. and Swenson, M.J. 1985. Minerals and Bones. In: Dukes' Physiology of Domestic Animals. 10th Ed. 449-466.
13. Jeejeebhoy KN (1999). Chromium and parenteral nutrition. *J. Trace Elem. Exp. Med.*, 2:85-89.
14. Lauwerys, R., Benard, A., Roelish., and Bouchet, J. P. 1994. Cadmium exposure markers as predictors of nephrotoxic effect. *Clin.*;40(7):1391-1394.
15. Mehra, R., and Juneja. M. 2003. Fingernails as Biological indices of metal exposure. *J. Biosci*; 30(2):253-257.
16. Merz, W. 1982. Clinical and public health significance of cadmium In : Clinical and biochemical nutritional aspects of trace elements. (Ed: As Prasad) Alan, R Liss, Inc., Newyork.
17. Nead, H. K., Halterman, J. S., and veteizman, M. 2004. Overweight children and adolescent: A risk group of iron deficiency. *Pediatric*; 114: 104-108.
18. Rodushkin I and Axelsson MD (2000) Application of double focusing sector field ICP-MS for multielemental characterization of human hair and nails. Part II. A study of the inhabitants of northern Sweden. *Sci Total Environ* 262:21–36
19. Shields GS, Coulson WF, Kimball DA, Cartwright GE, Wintrobe M (1962). Studies on copper metabolism XXXII: Cardiovascular lesions in copper-deficient swine. *Am. J. Pathol.*, 41: 603-621.
20. Tan, J.C., Burns, D.L. and Jones, H.R. 2006. Severe ataxia, myelopathy and peripheral neuropathy due to acquired copper deficiency in a patient with history of gastrectomy. *J. Paenteral. Nutr.* 30: 446-450.
21. Vincent JB (2004). Recent advances in the nutritional biochemistry of trivalent chromium. *Proc. Nutr. Soc.*, 63:41–7.
22. Were, F.H.; Njue, W.; Murungi, J. and Wanjau, 2008. *R. Sci. Total Environ*, 393, 376
